

CERTAIN SUBCLASS OF PASCU-TYPE BI-STARLIKE FUNCTIONS IN PARABOLIC DOMAIN

K. VIJAYA¹, §

ABSTRACT. Estimates on the coefficients $|a_2|$ and $|a_3|$ are obtained for normalized analytic function f in the open disk with f and its inverse $g = f^{-1}$ satisfy the condition that $\frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda z f'(z)}$ and $\frac{zg'(z) + \lambda z^2 g''(z)}{(1-\lambda)g(z) + \lambda z g'(z)}$ ($0 \leq \lambda \leq 1$) are both subordinate to an analytic function in parabolic region. Furthermore, we estimate the Fekete-Szegő functional for $f \in \mathcal{P}_{\Sigma, P}(\lambda, \varphi_\alpha)$.

Keywords: Analytic functions, univalent functions, bi-univalent functions, bi-starlike functions, bi-convex functions, and subordination.

AMS Subject Classification: 30C45.

1. INTRODUCTION

Let \mathcal{A} denote the class of analytic functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \tag{1}$$

normalized by the conditions $f(0) = 0 = f'(0) - 1$ defined in the open unit disk $\Delta = \{z \in \mathbb{C} : |z| < 1\}$. A function $f \in \mathcal{A}$ is said to be bi-univalent in Δ if both f and f^{-1} are univalent in Δ . Let Σ denote the class of bi-univalent functions defined in the unit disk Δ . Since $f \in \Sigma$ has the Maclaurian series given by (1), a computation shows that its inverse $g = f^{-1}$ has the expansion

$$g(w) = f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3)w^3 + \dots \tag{2}$$

An analytic function f is subordinate to an analytic function g , written $f(z) \prec g(z)$, provided there is an analytic function w defined on Δ with $w(0) = 0$ and $|w(z)| < 1$ satisfying $f(z) = g(w(z))$. Ma and Minda [8] unified various subclasses of starlike and convex functions for which either of the quantity $\frac{zf'(z)}{f(z)}$ or $1 + \frac{zf''(z)}{f'(z)}$ is subordinate to a more general superordinate function. For this purpose, they considered an analytic function φ with positive real part in the unit disk Δ , $\varphi(0) = 1, \varphi'(0) > 0$, and φ maps Δ onto a

¹ School of Advanced Sciences, Vellore Institute of Technology, Vellore-632 014, India.
e-mail: kvijaya@vit.ac.in; ORCID: <http://orcid.org/0000-0002-3216-7038>.

§ Manuscript received: April 10, 2109; accepted: May 5, 2020.

TWMS Journal of Applied and Engineering Mathematics, Vol.11, No.2 © Işık University, Department of Mathematics, 2021; all rights reserved.

region starlike with respect to 1 and symmetric with respect to the real axis. The class of Ma-Minda starlike functions consists of functions $f \in \mathcal{A}$ satisfying the subordination $\frac{zf'(z)}{f(z)} \prec \varphi(z)$. Similarly, the class of Ma-Minda convex functions of functions $f \in \mathcal{A}$ satisfying the subordination $1 + \frac{zf''(z)}{f'(z)} \prec \varphi(z)$. A function f is bi-starlike of Ma-Minda type or bi-convex of Ma-Minda type if both f and f^{-1} are respectively Ma-Minda starlike or convex. These classes are denoted respectively by $\mathcal{S}_{\Sigma}^*(\varphi)$ and $\mathcal{K}_{\Sigma}(\varphi)$. In the sequel, it is assumed that φ is an analytic function with positive real part in the unit disk Δ , satisfying $\varphi(0) = 1, \varphi'(0) > 0$, and $\varphi(\Delta)$ is symmetric with respect to the real axis. Such a function has a series expansion of the form

$$\varphi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + \dots, \quad (B_1 > 0). \tag{3}$$

Ali and Singh [2] introduced a new class of parabolic starlike functions denoted by $\mathcal{S}_p(\alpha)$ of order $\alpha (0 \leq \alpha < 1)$ satisfies the following:

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < (1 - 2\alpha) + \Re \left(\frac{zf'(z)}{f(z)} \right). \tag{4}$$

Equivalently,

$$f \in \mathcal{S}_p(\alpha) \iff \left(\frac{zf'(z)}{f(z)} \right) \in \Omega_{\alpha},$$

where Ω_{α} denotes the parabolic region in the right half-plane

$$\Omega_{\alpha} = \{w = u + iv : v^2 < 4(1 - \alpha)(u - \alpha)\} = \{w : |w - 1| < (1 - 2\alpha) + \Re(w)\}. \tag{5}$$

Ali and Singh [2] showed that the normalized Riemann mapping function $\varphi_{\alpha}(z)$ from the open unit disk Δ onto Ω_{α} is given by

$$\begin{aligned} \varphi_{\alpha}(z) &= 1 + \frac{4(1 - \alpha)}{\pi^2} \left[\log \frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right]^2 \\ &= 1 + \frac{16}{\pi^2}(1 - \alpha)z + \frac{32}{3\pi^2}(1 - \alpha)z^2 + \frac{368}{45\pi^2}(1 - \alpha)z^3 + \dots \\ &= 1 + \sum_{k=1}^{\infty} B_k z^k, \end{aligned} \tag{6}$$

where

$$B_k = \frac{16(1 - \alpha)}{k\pi^2} \sum_{j=0}^{k-1} \frac{1}{2j + 1} \quad (k \in \mathbb{N}). \tag{7}$$

Due to Ma and Minda [8], we state the following Lemma.

Lemma 1.1. *If a function $f \in \mathcal{S}_p(\alpha)$, then*

$$\left(\frac{zf'(z)}{f(z)} \right) \in \varphi_{\alpha}(z),$$

where φ_{α} is given by (6).

Since univalent functions are one-to-one, they are invertible and the inverse functions need not be defined on the entire unit disk Δ . In fact, the Koebe one-quarter theorem [6] ensures that the image of Δ . under every univalent function $f \in \mathcal{S}$ of the form (1), contains a disk of radius $\frac{1}{4}$. Thus every univalent function $f \in \mathcal{S}$ has an inverse f^{-1} which is defined by

$$f^{-1}(f(z)) = z \quad (z \in \Delta)$$

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f); r_0(f) \geq \frac{1}{4} \right).$$

In fact, the inverse function f^{-1} is given by

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots \quad (8)$$

Several authors have introduced and investigated subclasses of bi-univalent functions Σ and obtained bounds for the initial coefficients (see [4, 3, 10, 12]). Motivated by the work of Ali et al. [2, 7], in this paper, we introduce a new subclass $\mathcal{P}_{\Sigma,P}(\lambda, \varphi_\alpha)$ of bi-univalent functions and obtain the estimates on the coefficients $|a_2|$ and $|a_3|$ by subordination. Furthermore, we estimate the Fekete-Szegő functional for $f \in \mathcal{P}_{\Sigma,P}(\lambda, \varphi_\alpha)$.

Definition 1.1. A function $f \in \Sigma$ is said to be in the class $\mathcal{P}_{\Sigma,P}(\lambda, \varphi_\alpha)$ if the following subordination hold:

$$\left| \frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda z f'(z)} - 1 \right| < (1-2\alpha) + \Re \left(\frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda z f'(z)} \right) \quad (z \in \Delta) \quad (9)$$

and

$$\left| \frac{wg'(w) + \lambda w^2 g''(w)}{(1-\lambda)g(w) + \lambda w g'(w)} - 1 \right| < (1-2\alpha) + \Re \left(\frac{wg'(w) + \lambda w^2 g''(w)}{(1-\lambda)g(w) + \lambda w g'(w)} \right) \quad (w \in \Delta). \quad (10)$$

Due to Lemma 1.1 and by the above the definition we can state

$$\frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda z f'(z)} \prec \varphi_\alpha(z) \quad (z \in \Delta) \quad (11)$$

and

$$\frac{wg'(w) + \lambda w^2 g''(w)}{(1-\lambda)g(w) + \lambda w g'(w)} \prec \varphi_\alpha(w) \quad (w \in \Delta), \quad (12)$$

where φ_α is given by (6).

We note that $\mathcal{P}_{\Sigma,P}(0, \varphi_\alpha) = \mathcal{S}_{\Sigma,P}^*(\varphi_\alpha)$ [5] and $\mathcal{P}_{\Sigma,P}(1, \varphi_\alpha) = \mathcal{K}_{\Sigma,P}(\varphi_\alpha)$ as illustrated below:

Example 1.1. [5] A function $f \in \Sigma$ is said to be in the class $\mathcal{S}_{\Sigma,P}(\varphi_\alpha)$ if the following subordination hold:

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < (1-2\alpha) + \Re \left(\frac{zf'(z)}{f(z)} \right) \quad (z \in \Delta)$$

and

$$\left| \frac{wg'(w)}{g(w)} - 1 \right| < (1-2\alpha) + \Re \left(\frac{wg'(w)}{g(w)} \right) \quad (w \in \Delta).$$

Due to Lemma 1.1 and by the above the definition we can state

$$\frac{zf'(z)}{f(z)} \prec \varphi_\alpha(z) \quad \text{and} \quad \frac{wg'(w)}{g(w)} \prec \varphi_\alpha(w)$$

where $\varphi_\alpha(z)$ is given by (6) and $z, w \in \Delta$.

Example 1.2. A function $f \in \Sigma$ is said to be in the class $\mathcal{K}_{\Sigma,P}(\varphi_\alpha)$ if the following subordination hold:

$$\left| \frac{zf''(z)}{f'(z)} \right| < (1-2\alpha) + \Re \left(1 + \frac{zf''(z)}{f'(z)} \right) \quad (z \in \Delta)$$

and

$$\left| \frac{wg''(w)}{g'(w)} \right| < (1-2\alpha) + \Re \left(1 + \frac{wg''(w)}{g'(w)} \right) \quad (w \in \Delta).$$

Due to Lemma 1.1 and by the above the definition we can state

$$1 + \frac{zf''(z)}{f'(z)} \prec \varphi_\alpha(z) \quad \text{and} \quad 1 + \frac{wg''(w)}{g'(w)} \prec \varphi_\alpha(w)$$

where $\varphi_\alpha(z)$ is given by (6) and $z, w \in \Delta$. In order to prove our main results, we require the following Lemma due to [11].

Lemma 1.2. *If $h \in \mathcal{P}$, then $|c_k| \leq 2$ for each k , where \mathcal{P} is the family of all functions h analytic in Δ for which $\Re\{h(z)\} > 0$, where $h(z) = 1 + c_1z + c_2z^2 + \dots$ for $z \in \Delta$.*

2. SECTION

Coefficient estimates for the function class $\mathcal{P}_{\Sigma, P}(\lambda, \varphi_\alpha)$

Theorem 2.1. *Let f given by (1) be in the class $\mathcal{P}_{\Sigma, P}(\lambda, \varphi_\alpha)$. Then*

$$|a_2| \leq \frac{B_1\sqrt{B_1}}{\sqrt{|(1 + 2\lambda - \lambda^2)B_1^2 + (1 + \lambda)^2(B_1 - B_2)|}} \tag{13}$$

and

$$|a_3| \leq B_1 \left(\frac{B_1}{(1 + \lambda)^2} + \frac{1}{2(1 + 2\lambda)} \right). \tag{14}$$

where $B_1 = \frac{16}{\pi^2}(1 - \alpha)$ and $B_2 = \frac{32}{3\pi^2}(1 - \alpha)$ from (7).

Proof. Let $f \in \mathcal{P}_{\Sigma, P}(\lambda, \varphi_\alpha)$ and $g = f^{-1}$. Then there are analytic functions $u, v : \Delta \rightarrow \Delta$, with $u(0) = 0 = v(0)$, satisfying

$$\frac{zf'(z) + \lambda z^2 f''(z)}{(1 - \lambda)f(z) + \lambda z f'(z)} = \varphi_\alpha(u(z)) \tag{15}$$

and

$$\frac{wg'(w) + \lambda w^2 g''(w)}{(1 - \lambda)g(w) + \lambda w g'(w)} = \varphi_\alpha(v(w)). \tag{16}$$

Define the functions $p(z)$ and $q(z)$ by

$$p(z) := \frac{1 + u(z)}{1 - u(z)} = 1 + p_1z + p_2z^2 + \dots$$

and

$$q(z) := \frac{1 + v(z)}{1 - v(z)} = 1 + q_1z + q_2z^2 + \dots$$

or, equivalently,

$$u(z) := \frac{p(z) - 1}{p(z) + 1} = \frac{1}{2} \left[p_1z + \left(p_2 - \frac{p_1^2}{2} \right) z^2 + \dots \right] \tag{17}$$

and

$$v(z) := \frac{q(z) - 1}{q(z) + 1} = \frac{1}{2} \left[q_1z + \left(q_2 - \frac{q_1^2}{2} \right) z^2 + \dots \right]. \tag{18}$$

Then $p(z)$ and $q(z)$ are analytic in Δ with $p(0) = 1 = q(0)$. Since $u, v : \Delta \rightarrow \Delta$, the functions $p(z)$ and $q(z)$ have a positive real part in Δ , and $|p_i| \leq 2$ and $|q_i| \leq 2$. Using (17) and (18) in (15) and (16) respectively, we have

$$\frac{zf'(z) + \lambda z^2 f''(z)}{(1 - \lambda)f(z) + \lambda z f'(z)} = \varphi \left(\frac{1}{2} \left[p_1z + \left(p_2 - \frac{p_1^2}{2} \right) z^2 + \dots \right] \right) \tag{19}$$

and

$$\frac{wg'(w) + \lambda w^2 g''(w)}{(1-\lambda)g(w) + \lambda wg'(w)} = \varphi \left(\frac{1}{2} \left[q_1 w + \left(q_2 - \frac{q_1^2}{2} \right) w^2 + \dots \right] \right). \quad (20)$$

In light of (1) - (3), from (19) and (20), it is evident that

$$\begin{aligned} & 1 + (1 + \lambda)a_2 z + [2(1 + 2\lambda)a_3 - (1 + \lambda)^2 a_2^2] z^2 + \dots \\ &= 1 + \frac{1}{2} B_1 p_1 z + \left[\frac{1}{2} B_1 \left(p_2 - \frac{p_1^2}{2} \right) + \frac{1}{4} B_2 p_1^2 \right] z^2 + \dots \end{aligned}$$

and

$$\begin{aligned} & 1 - (1 + \lambda)a_2 w - [2(1 + 2\lambda)a_3 + (\lambda^2 - 6\lambda - 3)a_2^2] w^2 + \dots \\ &= 1 + \frac{1}{2} B_1 q_1 w + \left[\frac{1}{2} B_1 \left(q_2 - \frac{q_1^2}{2} \right) + \frac{1}{4} B_2 q_1^2 \right] w^2 + \dots \end{aligned}$$

which yields the following relations.

$$(1 + \lambda)a_2 = \frac{1}{2} B_1 p_1 \quad (21)$$

$$-(1 + \lambda)^2 a_2^2 + 2(1 + 2\lambda)a_3 = \frac{1}{2} B_1 \left(p_2 - \frac{p_1^2}{2} \right) + \frac{1}{4} B_2 p_1^2 \quad (22)$$

$$-(1 + \lambda)a_2 = \frac{1}{2} B_1 q_1 \quad (23)$$

and

$$-(\lambda^2 - 6\lambda - 3)a_2^2 - 2(1 + 2\lambda)a_3 = \frac{1}{2} B_1 \left(q_2 - \frac{q_1^2}{2} \right) + \frac{1}{4} B_2 q_1^2. \quad (24)$$

From (21) and (23), it follows that

$$p_1 = -q_1 \quad (25)$$

and

$$8(1 + \lambda)^2 a_2^2 = B_1^2 (p_1^2 + q_1^2). \quad (26)$$

From (22), (24) and (26), we obtain

$$a_2^2 = \frac{B_1^3 (p_2 + q_2)}{4[(1 + 2\lambda - \lambda^2)B_1^2 + (1 + \lambda)^2(B_1 - B_2)]}. \quad (27)$$

Applying Lemma 1.2, for the coefficients p_2 and q_2 , we immediately got the desired estimate on $|a_2|$ as asserted in (2.2).

By subtracting (24) from (22) and using (25) and (26), we get

$$a_3 = a_2^2 + \frac{B_1(p_2 - q_2)}{8(1 + 2\lambda)} = \frac{B_1^2(p_1^2 + q_1^2)}{8(1 + \lambda)^2} + \frac{B_1(p_2 - q_2)}{8(1 + 2\lambda)}. \quad (28)$$

Applying Lemma 1.2 once again for the coefficients p_1, p_2, q_1 and q_2 , we get the desired estimate on $|a_3|$ as asserted in (2.2) \square

Remark 2.1. For $\lambda = 0$ and $B_1 = \frac{16}{\pi^2}(1 - \alpha)$ and $B_2 = \frac{32}{3\pi^2}(1 - \alpha)$ the inequality (2.2) reduces to the estimate of $|a_2|$ and $|a_3|$. [5].

By taking $\lambda = 1$ we get the following result for $f \in \mathcal{K}_{\Sigma, P}(\varphi_\alpha)$

Theorem 2.2. Let f given by (1) be in the class $\mathcal{K}_{\Sigma, P}(\varphi_\alpha)$. Then

$$|a_2| \leq \frac{B_1 \sqrt{B_1}}{\sqrt{2B_1^2 + |4(B_1 - B_2)|}}$$

and

$$|a_3| \leq \frac{B_1^2}{4} + \frac{B_1}{6}.$$

where $B_1 = \frac{16}{\pi^2}(1 - \alpha)$ and $B_2 = \frac{32}{3\pi^2}(1 - \alpha)$ from (7).

2.1. Subsection. Fekete-Szegő inequalities for the Function Class $\mathcal{P}_{\Sigma,P}(\lambda, \varphi_\alpha)$ Making use of the values of a_2^2 and a_3 , and motivated by the recent work of Zaprawa [13], we prove the following Fekete-Szegő result for the function class $f \in \mathcal{P}_{\Sigma,P}(\lambda, \varphi_\alpha)$.

Theorem 2.3. *Let the function $f(z)$ be in the class $\mathcal{P}_{\Sigma,P}(\lambda, \varphi_\alpha)$ and $\mu \in \mathbb{C}$, then*

$$|a_3 - \mu a_2^2| \leq 2B_1 \left| \left(\Theta(\mu) + \frac{1}{8(1+2\lambda)} \right) + \left(\Theta(\mu) - \frac{1}{8(1+2\lambda)} \right) \right|, \tag{29}$$

where

$$\Theta(\mu) = \frac{B_1^2(1-\mu)}{4[(1+2\lambda-\lambda^2)B_1^2 + (1+\lambda)^2(B_1-B_2)]}, B_1 > 0.$$

where $B_1 = \frac{16}{\pi^2}(1 - \alpha)$ and $B_2 = \frac{32}{3\pi^2}(1 - \alpha)$ from (7)

Proof. From (28), we have

$$a_3 = a_2^2 + \frac{B_1(p_2 - q_2)}{8(1+2\lambda)}.$$

Using (27), by simple calculation we get

$$a_3 - \mu a_2^2 = B_1 \left[\left(\Theta(\mu) + \frac{1}{8(1+2\lambda)} \right) p_2 + \left(\Theta(\mu) - \frac{1}{8(1+2\lambda)} \right) q_2 \right],$$

where $\Theta(\mu) = \frac{B_1^2(1-\mu)}{4[(1+2\lambda-\lambda^2)B_1^2 + (1+\lambda)^2(B_1-B_2)]}$. Since all B_j are real and $B_1 > 0$, we have

$$|a_3 - \mu a_2^2| \leq 2B_1 \left| \left(\Theta(\mu) + \frac{1}{8(1+2\lambda)} \right) + \left(\Theta(\mu) - \frac{1}{8(1+2\lambda)} \right) \right|,$$

which completes the proof. □

Remark 2.2. *Specializing $\lambda = 0$ we can obtain the Fekete-Szegő inequality for the function class $\mathcal{S}_{\Sigma,P}(\varphi_\alpha)$ as in [5] .*

Specializing $\lambda = 1$ we can obtain the Fekete-Szegő inequality for the function class $\mathcal{K}_{\Sigma,P}(\varphi_\alpha)$ as given below.

Corollary 2.1. *Let the function $f(z)$ be in the class $\mathcal{K}_{\Sigma,P}(\varphi_\alpha)$ and $\mu \in \mathbb{C}$, then*

$$|a_3 - \mu a_2^2| \leq 2B_1 \left| \left(\Theta(\mu) + \frac{1}{24} \right) + \left(\Theta(\mu) - \frac{1}{24} \right) \right|,$$

where

$$\Theta(\mu) = \frac{B_1^2(1-\mu)}{4[2B_1^2 + 4(B_1 - B_2)]}, B_1 > 0.$$

where $B_1 = \frac{16}{\pi^2}(1 - \alpha)$ and $B_2 = \frac{32}{3\pi^2}(1 - \alpha)$ from (7)

3. CONCLUSIONS

By taking $B_1 = \frac{16}{\pi^2}(1 - \alpha)$ and $B_2 = \frac{32}{3\pi^2}(1 - \alpha)$ and specializing the parameter $\lambda = 1$ we state the results for the class of bi convex functions in parabolic domain which has not been studied. Further by specializing $\lambda = 0$ we can obtain the results for bi-starlike functions in parabolic domain as in [5] .

REFERENCES

- [1] Ali R.M.and Singh V. , (1994), Coefficients of parabolic starlike functions of order α , Computational Methods and Function Theory(Penang), Ser. Approx. Decompos., Vol.5, pp.23–36. World Scientific Publishing, New Jersey(1995).
- [2] Ali R.M., Leo S.K., Ravichandran V., Supramaniam S.,(2012), Coefficient estimates for bi-univalent Ma-Minda star-like and convex functions, Appl. Math. Lett. 25, pp344 -351.
- [3] Brannan D.A., Clunie J.,(1970), W.E. Kirwan, Coefficient estimates for a class of star-like functions, Canad. J. Math. 22, pp476 - 485.
- [4] Brannan D.A., Taha T.S.,(1986), On some classes of bi-univalent functions, Studia Univ. Babeş-Bolyai Math. 31(2), pp 70 - 77.
- [5] Bulut S.,(2018), *Coefficient estimates for a subclass of parabolic bi-starlike functions*, Afr. Mat. **29** Issue 34, pp 331–338.
- [6] Duren P.L.,(1983), Univalent Functions, in: Grundlehren der Mathematischen Wissenschaften, Vol. 259, Springer, New York.
- [7] Lewin M.,(1967) On a coefficient problem for bi-univalent functions, Proc. Amer. Math. Soc. 18, pp 63 - 68.
- [8] Ma W.C., Minda D.,(1992), A unified treatment of some special classes of functions, in: Proceedings of the Conference on Complex Analysis, Tianjin,, pp 157 - 169, Conf. Proc. Lecture Notes Anal. 1. Int. Press, Cambridge, MA, 1994.
- [9] Netanyahu E.,(1969), The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$. Arch. Ration. Mech. Anal. 32 pp 100 -112.
- [10] Murugusundaramoorthy, G., Cho, N.E.,(2019) On λ - pseudo bi-starlike functions in parabolic domain, Nonlinear Functional Analysis and Applications, 24(1), pp. 185-194
- [11] Pommerenke Ch.,(1975), Univalent functions, Vandenhoeck and Ruprecht, Göttingen,.
- [12] Srivastava H.M., Mishra A.K., Gochhayat P.,(2010), Certain subclasses of analytic and bi-univalent functions, Appl. Math. Lett. 23(10) pp 1188 - 1192.
- [13] Zaprawa P. ,(2014), On the Fekete-Szegő problem for classes of bi-univalent functions, Bull. Belg. Math. Soc. Simon Stevin 21(1), pp1–192.
- [14] Erdelyi, A., (1956), Asymptotic expansions, Dover publications, New York.
- [15] Anderson, J. D., Campbell, J. K., Ekelund, J. E., Ellis, J. and Jordan, J. F., (2008), Physical Review Letters, 100, 091102.
- [16] Mould, R. A., (1994), Basic Relativity, Springer-Verlag Newyork Inc.
- [17] Thompson, K. W., (1987), Time dependent boundary conditions for hyperbolic systems, J. Comp. Phys., 68, pp. 1-24.
- [18] Hixon, R. and Turkel, E., (2000), Compact implicit MacCormack-type schemes with high accuracy, J. Comp. Phys., 158, pp. 51-70.
- [19] Fan, E. and Jian, Z., (2002), Applications of the Jacobi elliptic function method to special-type nonlinear equations, Phys. Lett. A, 305 (6), pp. 383-392.

Vijaya Kaliyappan for the photography and short autobiography, see TWMS J. App. and Eng. Math., V.11, N.2, 2021, p.469-479
